



# A Pharmacology Perspective on Simultaneous Tuberculosis and Hepatitis C Treatment

 Russell R. Kempker,<sup>a</sup>  Wael A. Alghamdi,<sup>b</sup>  Mohammad H. Al-Shaer,<sup>c</sup>  Gena Burch,<sup>c</sup>  Charles A. Peloquin<sup>c</sup>

<sup>a</sup>Division of Infectious Diseases, Department of Medicine, Emory University School of Medicine, Atlanta, Georgia, USA

<sup>b</sup>Department of Clinical Pharmacy, College of Pharmacy, King Khalid University, Abha, Saudi Arabia

<sup>c</sup>Department of Pharmacotherapy and Translational Research, College of Pharmacy, University of Florida, Gainesville, Florida, USA

**ABSTRACT** Tuberculosis (TB) and hepatitis C virus (HCV) infections are both major public health problems. Despite high rates of coinfection, there is scarce literature addressing the convergence of the two diseases. One particularly unexplored area is the potential for simultaneous treatment of TB and HCV which would allow for leveraging an extensive global TB treatment infrastructure to help scale up HCV treatment. We review the drug metabolism of anti-TB and HCV drugs and the known and potential drug-drug interactions between recommended HCV regimens and individual anti-TB drugs. Rifampin is the only anti-TB drug to have been formally studied for potential drug interactions with anti-HCV direct-acting antivirals (DAAs), and existing data preclude these combinations. However, based on known pathways of drug metabolism and enzyme effects, the combination of HCV DAA regimens with all other anti-TB drugs may be feasible. Pharmacokinetic studies are needed next to help move cotreatment regimens forward for clinical use among patients coinfecting with TB and HCV.

**KEYWORDS** pharmacology, tuberculosis, hepatitis C, drug interactions, cotreatment

Tuberculosis (TB) and chronic hepatitis C virus (HCV) infections are both major public health problems responsible for significant morbidity and mortality worldwide. There are an estimated 10 million incidence cases of TB globally and approximately 1.6 million TB-related deaths per year, making it the leading cause of infectious disease-related mortality worldwide (1). Infection with HCV leads to chronic hepatitis in the majority of cases, and in addition to >70 million prevalent cases, there are an estimated 3 to 4 million new cases and 400,000 deaths annually (2). Disease burden for both TB and HCV are high and disproportionately affect persons in low- and middle-income countries (LMICs) (1, 2). While data are limited, the prevalence of HCV has been found to be high (>5%) among patients with active TB, especially in Eastern Europe and Central Asia (>15%), where high rates of incarceration and injection drug use fuel both epidemics (3, 4). Beyond studies focusing on the coprevalence of TB and HCV (5) or rates of hepatotoxicity (6–8), there has been scarce discussion on the potential synergies of integrating TB and HCV treatment.

The development and rollout of new all oral direct-acting antivirals (DAAs) for the treatment of HCV infection have occurred at breakneck speed and have revolutionized HCV treatment, leading to extremely high cure rates and improved quality of life among those treated (9, 10). However, there are many challenges to expanding HCV treatment with DAAs, including improving case finding for chronic HCV infection (~20% of people with chronic HCV infection know their status), high drug costs, and the development of a health care workforce and infrastructure to provide treatment (2). In 2015, it was estimated that only 7% of people diagnosed with chronic HCV infection were started on treatment and access to DAAs, especially in most LMICs, remains

**Citation** Kempker RR, Alghamdi WA, Al-Shaer MH, Burch G, Peloquin CA. 2019. A pharmacology perspective on simultaneous tuberculosis and hepatitis C treatment. *Antimicrob Agents Chemother* 63:e01215-19. <https://doi.org/10.1128/AAC.01215-19>.

**Copyright** © 2019 American Society for Microbiology. All Rights Reserved.

Address correspondence to Russell R. Kempker, [rkempke@emory.edu](mailto:rkempke@emory.edu).

**Accepted manuscript posted online** 7 October 2019

**Published** 21 November 2019

limited (2). Given the lack of capacity to treat HCV infection in many settings, including in both high-income countries and LMICs, there have been efforts and discussion to leverage existing primary care, HIV, and injection drug use (IDU) treatment programs to administer HCV treatment (11). An additional unexplored area is the use of TB treatment facilities to provide HCV treatment for TB/HCV coinfecting patients. National TB programs exist in most countries, and their experienced and skilled health care workforce consisting of physicians and nurses along with extensive associated laboratory networks could offer a vital resource for HCV management in areas with insufficient capacity to meet increasing HCV treatment demand. Treating TB/HCV coinfecting patients at a TB facility would also take advantage of the existing therapeutic relationship between TB health care workers and their patients, which develops throughout the long duration of TB treatment and would be in line with the goal of providing patient-centered care to all patients with TB.

To consider the cotreatment of patients with active TB and chronic HCV infection, it is imperative to understand the pharmacology of anti-TB and HCV drugs and consider possible drug-drug interactions (DDIs) that may allow or preclude the use of certain drug combinations. To our knowledge, there has been only one report in the literature of a patient simultaneously treated for active TB and HCV (12) and one published study on anti-TB and anti-HCV DDIs in healthy subjects (13). To evaluate the pharmacology of simultaneous TB and HCV treatment, we first briefly review the metabolism of anti-TB drugs and oral DAAs for HCV and then discuss potential DDIs; lastly, we discuss other issues related to the pharmacology of TB and HCV cotreatment, including liver disease and drug use. Our goal is to drive the field of TB and HCV comanagement ahead and ultimately improve treatment outcomes by providing data to help guide a research agenda for TB and HCV pharmacokinetic and clinical treatment studies.

## TB AND HCV DRUG METABOLISM

For anti-TB drugs, we review the metabolism of drugs recommended for drug-susceptible (DS) and multidrug-resistant (MDR) TB (14–16). With regard to HCV drugs, we cover the metabolism of drugs that are currently recommended as either first-line or alternative treatment regimens by the American Association for the Study of Liver Diseases (AASLD) and the Infectious Diseases Society of America (IDSA) (17). For a more exhaustive review, we direct the reader to recent in depth reviews (18–20). An overview of key routes of metabolism for anti-TB and HCV drugs are displayed in Tables 1 and 2, respectively.

**Anti-TB drugs.** The rifamycins are a cornerstone of first-line anti-TB treatment, and they also present the most challenging drug class in regard to DDIs, given their various and potent effects on human drug metabolism systems (21). To a great extent, rifamycins are eliminated by intestinal and hepatic metabolism to mostly desacetylated and hydroxylated metabolites (22). The three rifamycins used to treat mycobacterial infections include rifampin, rifapentine, and rifabutin, and all are metabolized to a partially active 25-*O*-desacetyl metabolite. Rifampin and rifapentine are cleared largely by an esterase, whereas rifabutin and especially 25-*O*-desacetyl-rifabutin depend on cytochrome P450 (CYP) 3A (CYP3A) for enzymatic clearance. Both the rifamycins and their metabolites are excreted primarily in the bile and eliminated in the feces (22). Approximately 10% of rifabutin, 17% of rifapentine, and 13% to 24% of rifampin are excreted unchanged in the urine (23–25).

Rifampin has an elimination half-life ( $t_{1/2}$ ) of approximately 2 to 4 h (24). Rifampin notably induces its own metabolism or intestinal efflux (autoinduction), with the result that area under the concentration-time curve (AUC) and plasma  $t_{1/2}$  are both reduced by 20% to 40% after 7 to 10 days of daily dosing (26). This autoinduction process is different from rifampin's well-known effects on CYP enzymes, since rifampin is not a substrate for CYP enzymes (24, 27). Rifampin causes increased metabolism of many drugs, but it rarely is affected by other drugs. Some genetic variations in carboxylesterase 2 enzyme may alter rifampin metabolism by affecting the gene expression (28).

**TABLE 1** Elimination of antituberculosis drugs and their effects on enzyme and transporter activity<sup>a</sup>

Drug	Reference(s)	Route of Elimination	Transporter substrate	Enzyme/transporter inhibition	Enzyme/transporter induction
<b>First-line drugs</b>					
Rifampin	22, 24	Deacetylation	OATP1B1/SLCO1B1, P-gp/ABCB1	Transporters: OATP1B1/SLCO1B1	Enzymes: CYP1A2, -2B6, -2C19, -2C8, -2C9, -3A4, UGT1A1. Transporters: P-gp/ABCB1
Rifabutin	22, 34	CYP3A, hydroxylation, deacetylation	—	—	CYP3A4
Rifapentine	22, 35, 39, 41	Deacetylation, hydrolysis	—	—	CYP3A4, -2C8/9
Isoniazid	43, 45	Acetylation, dehydrazination	—	Enzymes: CYP2C19, -3A, -2A6, -2E1	CYP2E1
Pyrazinamide	46	Hydrolysis	—	—	—
Ethambutol	47	20% oxidation, 80% excreted unchanged in urine	—	—	—
<b>Second-line drugs</b>					
Levofloxacin	49	87% excreted unchanged in urine	—	—	—
Moxifloxacin	55, 56	Glucuronide and sulfate conjugation	P-gp	—	—
Amikacin	47	>90% excreted unchanged in urine	—	—	—
Kanamycin	47	>90% excreted unchanged in urine	—	—	—
Capreomycin	47	50–60% excreted renally	—	—	—
Ethionamide	51	Oxidation to sulphoxide metabolite	—	Enzyme: CYP2C8	—
Cycloserine	47	50–70% excreted unchanged in urine	—	—	—
Linezolid	50	Metabolized by oxidation (independent of CYP enzymes); renal and non-renal excretion	P-gp	—	—
Clofazimine	57, 93	Hydrolysis and glucuronidation	—	Enzymes: CYP3A, P-gp, BCRP, MRP1	—
Bedaquiline	58	CYP3A4 (mainly), CYP2C8, CYP2C19	—	—	—
Delamanid	59, 60	Albumin; metabolites are further metabolized by CYP enzymes (mainly by CYP3A4)	—	—	—
PAS	53	Acetylation	—	—	—

<sup>a</sup>—, either unknown or no well-established effect. ABC, ATP-binding cassette transporters; CYP, cytochrome 450; OATP, organic-anion-transporting polypeptide; BCRP, breast cancer resistance protein; MRP1, multidrug resistance-associated protein 1; P-gp, P-glycoprotein; PAS, *p*-aminosalicylic acid; SLC, solute carrier family; UGT, uridine 5'-diphospho-glucuronosyltransferase.

Rifabutin undergoes extensive intestinal and hepatic metabolism, resulting in more than 20 metabolites (22, 29). The primary metabolites are 25-*O*-desacetyl-rifabutin and 31-hydroxyl-rifabutin (22, 23, 29, 30). Rifabutin elimination  $t_{1/2}$  is in the range of 25 to 67 h, depending on whether the data are obtained using a single dose (longer) or at steady state (shorter) (22, 23, 31). Similarly to rifampin, rifabutin undergoes autoinduction with 37% and 13% decreases in AUC and maximum concentration ( $C_{max}$ ), respectively, after 10 daily doses (22, 29, 32, 33). CYP3A is the major isozyme that transforms rifabutin to its oxidative metabolites in human enterocytes and liver microsomes, and it also catalyzes oxidation of 25-*O*-desacetyl-rifabutin (22, 29). Rifabutin excretion occurs through the biliary tract and, to a small degree, renally (30). After a 150-mg dose, biliary concentrations reach up to 3 to 5 times the plasma concentration (34). Although not free of enzyme induction, rifabutin typically is approximately 40% as potent as rifampin and has been the preferred rifamycin when DDIs are unavoidable (34).

The elimination  $t_{1/2}$  of rifapentine ranges from 14 to 18 h in healthy young adults to ~20 h in men >65 years (35) and is reduced with repeated dosing, suggesting autoinduction (36, 37). Rifapentine is metabolized by arylacetamide deacetylase (38), with nearly all radiolabeled drug in plasma being accounted for by rifapentine and its 25-desacetyl metabolite (39, 40). Approximately 87% of radiolabeled rifapentine is recovered in either feces (70%) or urine (17%) (41).

With regard to the remaining first-line drugs, isoniazid is extensively metabolized, mainly in the liver, to a number of inactive compounds by acetylation and dehydrazination. *N*-Acetyltransferase 2 forms acetyl-isoniazid, which is further metabolized to mono- and diacetylhydrazine (42–44). Isoniazid is an inhibitor of CYP2A6, -3A, -2C19, and -2E1, which may interact with other drugs with a narrow therapeutic index (45). Pyrazinamide undergoes hydrolysis (46), and the majority of ethambutol is excreted unchanged in the urine (47).

**TABLE 2** Elimination of DAAs for hepatitis C and their effects on enzymes and transporters activity<sup>a</sup>

Drug	Reference(s)	Route of Elimination	Transporter substrate	Enzyme/transporter inhibition	Enzyme/transporter induction
Elbasvir	69	CYP3A4	P-gp	Transporters: BCRP, P-gp	None
Daclatasvir	68	CYP3A	P-gp	Enzymes: CYP3A4. Transporters: P-gp, BCRP, OATP1B1/3	CYP3A4 (weak)
Glecaprevir	70	CYP3A4 (secondary)	P-gp, BCRP, OATP1B1/3	Enzymes: CYP3A, CYP1A2, UGT1A1. Transporters: P-gp, BCRP, OATP1B1/3	—
Grazoprevir	69	CYP3A4	P-gp, OATP1B1	Enzymes: CYP3A4, UGT1A1. Transporter: BCRP	—
Ledipasvir	65, 66	Oxidative metabolism via unknown route	P-gp, BCRP	Enzymes: intestinal CYP3A4, UGT. Transporters: P-gp, BCRP	CYP3A4 (weak), CYP2C (weak), UGT1A1 (weak)
Pibrentasvir	70	None; eliminated via biliary-fecal route	P-gp, BCRP	Enzymes: CYP3A, -1A2, UGT1A1. Transporters: P-gp, BCRP, OATP1B1/3	—
Sofosbuvir	62, 66	Phosphorylated, renal excretion	P-gp, BCRP	—	—
Velpatasvir	72	CYP2B6, CYP2C8, CYP3A4	P-gp, BCRP	Transporters: P-gp, BCRP, OATP1B1/3, OATP2B1	—
Voxilaprevir	71	Extensive via CYP3A4	P-gp, BCRP, OAT1B1/3	Transporters: P-gp, BCRP, OATP1B1/3	—
Paritaprevir	67	CYP3A4, CYP3A5 (minor)	P-gp, BCRP, OAT1B1/3	Enzyme: UGT1A1. Transporters: BCRP, OATP1B1/3	—
Ombitasvir	67	Amide hydrolysis and oxidative metabolism	P-gp, BCRP	Enzyme: UGT1A1	—
Dasabuvir	67	CYP2C8, CYP3A4 (minor)	P-gp, BCRP	Enzyme: UGT1A1. Transporter: BCRP	—
Simeprevir	64	CYP3A4, possibly CYP2C8, CYP2C19	P-gp, OATP1B1/3, OATP2B1	Enzymes: intestinal CYP3A4 (mild), CYP1A2 (mild). Transporters: P-gp, OATP1B1/3	—

<sup>a</sup>—, either unknown or no well-established effect. BCRP, breast cancer resistance protein; CYP, cytochrome 450; DAAs, direct-acting antiviral; OATP, organic-anion-transporting polypeptide; P-gp, P-glycoprotein; UGT, uridine 5'-diphospho-glucuronosyltransferase.

Many of the second-line anti-TB drugs appear to be metabolized by other non-CYP routes and thus are not affected by CYP inducers or inhibitors; hence, fewer DDIs are expected overall. Aminoglycosides, capreomycin, levofloxacin, and cycloserine are mostly excreted unchanged in the urine (47–49). Ethionamide and linezolid are mainly metabolized by oxidation (50, 51), and *p*-aminosalicylic acid undergoes acetylation (52, 53). Finally, moxifloxacin and clofazimine are metabolized mainly by conjugation (54, 55). Moxifloxacin is a P-glycoprotein (P-gp) substrate (56), while clofazimine is a P-gp inhibitor (57).

Bedaquiline and delamanid are the first new drugs to be approved for TB in >40 years, and they have brought much hope and promise to improve the management of MDR-TB. Bedaquiline is eliminated hepatically, being metabolized predominantly by CYP3A4 to its *N*-desmethyl metabolite. CYP2C8 and -2C19 have also been found to contribute to its metabolism (58). Delamanid, on the other hand, has a unique metabolic pathway. The parent drug is metabolized by plasma albumin to DM-6705 (M1), which is further metabolized by CYP enzymes through multiple metabolic pathways to form several metabolites (M2 to M8) (59, 60). The predominant pathway is mediated by CYP3A4 to form its oxide (M2).

**Anti-HCV drugs.** The route and extent of metabolism is varied among the oral DAAs (Table 2). Boceprevir and telaprevir were the first DAAs to be introduced and are metabolized by CYP3A4 and are P-gp substrates; consequently, they are prone to many DDIs (61). Given decreasing demand and availability of improved HCV treatments, both boceprevir and telaprevir have been pulled from the market. More recently introduced DAAs, sofosbuvir and simeprevir, are extensively metabolized by the liver. Sofosbuvir is a phosphoramidate prodrug that is hydrolyzed by intracellular cathepsin A and/or carboxylesterase 1 to the inactive metabolite GS-331007 (monophosphate), which is sequentially phosphorylated to the active metabolite GS-461203 (triphosphate) (62, 63). Simeprevir is known to be primarily metabolized by CYP3A4, with possible additional involvement of CYP2C8 and CYP2C19 (64).

The agents formulated in the multiple combination regimens approved between the years 2014 and 2017 also exhibit variable metabolism. Ledipasvir undergoes minimal oxidative metabolism through an unknown route and also requires an acidic pH for absorption (65, 66). Ombitasvir undergoes amide hydrolysis followed by oxidative metabolism in various locations (67). Paritaprevir is predominantly metabolized by CYP3A4 and secondarily through CYP3A5. Daclatasvir, elbasvir, and grazoprevir are also metabolized by CYP3A4 (68, 69). In contrast, dasabuvir is mainly metabolized by CYP2C8, while CYP3A4 is a minor pathway of metabolism. Velpatasvir undergoes extensive liver metabolism and has the potential to be implicated in DDIs due to the fact that it requires an acidic gastric environment for absorption and is metabolized by multiple CYP enzymes, including CYP2B6, -3A4, and -2C8 (65). Glecaprevir is secondarily metabolized by CYP3A4, but pibrentasvir does not undergo any metabolism. Both are primarily eliminated through the biliary-fecal route (70). The most recent agent approved in combination, voxilaprevir, is extensively metabolized in the liver by CYP3A4 (71).

### DDIs BETWEEN ANTI-TB AND HCV DRUGS

As there are currently limited published DDI studies of anti-TB drugs and HCV DAAs, we used data from drug package inserts and knowledge of drug metabolism to determine known and potential DDIs, respectively. With regard to available DDI studies contained in drug inserts, the only anti-TB drug studied was rifampin; thus, we extrapolated potential DDIs for the remainder of anti-TB and HCV DAA drug combinations. To further evaluate and confirm potential DDIs, we also utilized the University of Liverpool HEP drug interaction online tool (<https://www.hep-druginteractions.org>), with the exception of the following anti-TB drugs, which were not included in this database: kanamycin, cycloserine, clofazimine, ethionamide, and *para*-aminosalicylic acid. We evaluated potential DDIs from the perspective of different HCV regimens and

**TABLE 3** Drug-drug interactions between drug-susceptible tuberculosis and hepatitis C drugs<sup>a</sup>

Hepatitis C Treatment Regimens* (genotypes covered)	Anti-Tuberculosis Drugs for Drug-Susceptible Disease <sup>#</sup>					
	INH	RIF	RFP	RFB	PZA	ETH
First-line						
Glecaprevir, pibrentasvir (1-6)						
Sofosbuvir, velpatasvir (1-6)						
Elbasvir, grazoprevir (1,4)						
Ledipasvir, sofosbuvir (1,4-6)						
Alternative						
Paritaprevir, ritonavir, ombitasvir, dasabuvir, ribavirin (1)						
Paritaprevir, ritonavir, ombitasvir, ribavirin (4)						
Simeprevir, sofosbuvir (1)						
Daclatasvir, sofosbuvir (1-3)						
Elbasvir, grazoprevir, ribavirin (1)						

<sup>a</sup>\*, as recommended in the AALSD/IDSA hepatitis C treatment guidelines; #, drugs put into drug-susceptible and drug-resistant categories based on where they are commonly utilized (overlap does exist); INH, isoniazid; RIF, rifampin; RFP, rifapentine; RFB, rifabutin; PZA, pyrazinamide; ETH, ethambutol; red, drugs that should not be coadministered; orange, potential clinically significant interaction; yellow, potential weak interaction unlikely to be clinically significant; green, no clinically significant interaction expected.

their predicted interaction with individualized anti-TB drugs used for DS- and MDR-TB as shown in Tables 3 and 4.

**First-line HCV regimens.** Currently, there are two first-line pangenotypic drug regimens for the treatment of HCV infection, including glecaprevir-pibrentasvir and

**TABLE 4** Drug-drug interactions between drug-resistant tuberculosis and hepatitis C drugs<sup>a</sup>

Hepatitis C Treatment Regimens* (genotypes covered)	Anti-Tuberculosis Drugs for Drug-Resistant Disease <sup>#</sup>						
	LEV MOX	AMK KAN CAP	LZD	CFZ	PAS CS ETN	BDQ	DEL
First-line							
Glecaprevir, pibrentasvir (1-6)							
Sofosbuvir, velpatasvir (1-6)							
Elbasvir, grazoprevir (1,4)(118)							
Ledipasvir, sofosbuvir (1,4-6)							
Alternative							
Paritaprevir, ritonavir, ombitasvir, dasabuvir, ribavirin (1)							
Paritaprevir, ritonavir, ombitasvir, ribavirin (4)							
Simeprevir, sofosbuvir (1)							
Daclatasvir, sofosbuvir (1-3)							
Elbasvir, grazoprevir, ribavirin (1)							

<sup>a</sup>\*, as recommended in the AALSD/IDSA hepatitis C treatment guidelines; #, drugs put into drug-susceptible and drug-resistant categories based on where they are commonly utilized (overlap does exist); LEVO, levofloxacin; MOXI, moxifloxacin; AMK, amikacin; KAN, kanamycin; CAP, capreomycin; LZD, linezolid; CFZ, cefazolin; PAS, *para*-aminosalicylic acid; CS, cycloserine; ETN, ethionamide; BDQ, bedaquiline; DEL, delamanid; orange, potential clinically significant interaction; yellow, potential weak interaction unlikely to be clinically significant; green, no clinically significant interaction expected. Genotypes for first-line treatment with elbasvir and grazoprevir, see reference 94.



sofosbuvir-velpatasvir, and two additional regimens available for specific HCV genotypes, including elbasvir-grazoprevir and sofosbuvir-ledipasvir. Potential DDIs are reviewed below.

Glecaprevir and pibrentasvir are substrates and inhibitors of P-gp and breast cancer resistance protein (BCRP), with glecaprevir also being a substrate and inhibitor of organic-anion-transporting polypeptide (OATP) 1B1/3 (70). Additionally, both drugs are weak inhibitors of CYP3A, CYP1A2, and uridine 5'-diphospho-glucuronosyltransferase (UGT) 1A1. When given with 600 mg rifampin once daily, the  $C_{max}$  and AUC of both glecaprevir and pibrentasvir were decreased >80%; hence, rifampin is not recommended to be given with glecaprevir-pibrentasvir (70). Rifabutin has been used successfully with anti-HIV drugs and should be tested with anti-HCV drugs. Based on existing knowledge of drug metabolism, other potentially clinically significant DDIs include bedaquiline, which is metabolized by CYP3A4; thus, its concentrations may be increased with glecaprevir-pibrentasvir. Weaker potential DDIs are possible with isoniazid or clofazimine given their inhibitory effect on CYP3A enzymes which are responsible in part for the metabolism of glecaprevir. Clofazimine also inhibits two of the transporters (BCRP and P-gp) that glecaprevir and pibrentasvir are substrates for (57).

With regard to sofosbuvir and velpatasvir, both are substrates of P-gp and BCRP transporters. Additionally, velpatasvir is slowly metabolized by various CYP enzymes and is an inhibitor of BCRP and OATP transporters. The  $C_{max}$  and AUC of both sofosbuvir and velpatasvir were significantly decreased (>70%) in the presence of continuous daily use of rifampin 600 mg; thus, combination use with rifampin and other rifamycins currently is contraindicated (72). Additional potential DDIs exist between isoniazid and clofazimine and velpatasvir given their weak inhibition of CYP3A enzymes. There are no other predicted significant interactions with additional anti-TB drugs given the lack of any induction or inhibition of CYP enzymes by sofosbuvir or velpatasvir. Of note, the DDIs with the salvage regimen of sofosbuvir-velpatasvir-voxilaprevir and anti-TB medications would be expected to be similar to those described above, given both velpatasvir and voxilaprevir are eliminated by CYP enzymes (71).

The combination of elbasvir and grazoprevir is approved for HCV genotypes 1 to 4; both drugs are metabolized by CYP3A4 and are substrates of P-gp. Grazoprevir is also a substrate of OATP1B1. Both drugs can inhibit BCRP. Elbasvir is a weak inhibitor of P-gp while grazoprevir is a weak inhibitor of CYP3A4 and UGT1A1. While elbasvir-grazoprevir is not recommended to be given with rifampin, DDI data demonstrate mixed effects, with single-dose 600 mg rifampin increasing the AUC and  $C_{max}$  of elbasvir and grazoprevir, and continuous 600-mg-daily rifampin having minimal effect on grazoprevir AUC and  $C_{max}$  and decreasing 24-h postdrug concentrations by 90% (69). The inhibition of CYP3A4 by grazoprevir may increase the concentrations of bedaquiline, and given the uncertainties of the therapeutic index of bedaquiline, this is a potentially clinically significant interaction. Isoniazid and clofazimine may increase concentrations of both elbasvir and grazoprevir given their weak inhibition of CYP3A enzymes. Other anti-TB drugs are expected to have no major DDIs with elbasvir-grazoprevir.

The combination of sofosbuvir and ledipasvir has been the most commonly used DAA combination to date and is approved for genotypes 1 and 4 to 6. Similar to sofosbuvir, ledipasvir is also a substrate of P-gp, but in contrast, is also an inhibitor of P-gp and BCRP. A single dose of 600 mg rifampin decreased the  $C_{max}$  and AUC of ledipasvir by 35% and 59%, respectively. With regard to the clinical impact of this decrease, initial dose-finding studies found little additional antiviral activity with ledipasvir >30 mg; however, viral breakthrough with use of 30 mg compared to the standard 90-mg dose in one study was almost double (20% versus 11%) (73, 74). While the exposure-response relationships of sofosbuvir and ledipasvir need to be better defined, the pharmacokinetic impact of rifampin on ledipasvir along with effects on sofosbuvir mentioned above currently precludes the use of sofosbuvir-ledipasvir with rifampin or other rifamycins in clinical practice (66). Given low to no effect on the CYP enzyme system, there is no other clinically significant interaction with additional anti-TB drugs.

**Alternative recommended HCV regimens.** There are an additional five combination regimens recommended as alternative treatment regimens for HCV genotypes 1 to 4 as outlined in Table 3. With regard to the elbasvir-grazoprevir plus ribavirin combination for genotype 1, the expected DDIs are similar to the elbasvir-grazoprevir mentioned in the above section given no likely DDIs between ribavirin and anti-TB medications. The DDIs for the remaining 4 regimens are discussed below.

For the two regimens containing sofosbuvir, including simeprevir-sofosbuvir and daclatasvir-sofosbuvir, there are potential DDIs due to the metabolism of simeprevir and daclatasvir by CYP3A4 and the inhibition of CYP3A4 by daclatasvir and transporters by both drugs. Daily 600-mg rifampin decreased the  $C_{\max}$  and AUC of daclatasvir by 79% and 56% (68), respectively, while it increased the  $C_{\max}$  of simeprevir by 31% but decreased its AUC by 48% and  $C_{\min}$  by 92% (64). Hence, rifampin and all rifamycins currently are not recommended to be coadministered with either of these regimens. Given a mild inhibition of CYP3A4 by simeprevir, there is a potential increase in bedaquiline concentrations which warrants caution. Conversely, there is no expected interaction between daclatasvir and bedaquiline.

The two regimens based on a backbone of paritaprevir, ritonavir, ombitasvir, and ribavirin plus/minus dasabuvir have no available data including rifampin but have a high potential for DDIs (67). Both paritaprevir and dasabuvir are metabolized by CYP enzymes, which likely prohibit their use with the CYP-inducing rifamycins. Additionally, ritonavir is a strong CYP3A4, which raises the possibility of DDIs with anti-TB drugs, particularly bedaquiline and delamanid. When administered with lopinavir-ritonavir, the AUC of delamanid and its metabolite (DM-6705) was found to be increased by 25% (13), and the clearance of bedaquiline was decreased by 35%, leading to an estimated 3-fold increase in exposure during chronic treatment (75). The inhibition of UGT1A1 by paritaprevir, ombitasvir, and dasabuvir may also cause increased drug concentrations of drugs that undergo glucuronidation, including moxifloxacin.

**Overview.** We conclude with a few key points from our review of the literature. The use of rifampin with recommended HCV DAA regimens is contraindicated given the strong induction of numerous drug-metabolizing systems and given available data provided through pharmaceutical studies conducted. This would preclude the cotreatment of HCV infection with DAAs and drug-susceptible TB with current first-line rifampin-based regimens. Further pharmacokinetic study of elbasvir-grazoprevir with rifampin and DDI studies of DAAs with rifamycins that have less drug-metabolizing induction effects, specifically, rifabutin, are possible areas of further study. Rifabutin is a potentially attractive rifamycin to use with ant-HCV drugs given its lower potency of CYP enzyme induction than with rifampin; given that it may be feasible to use rifabutin with DAAs from a pharmacokinetic perspective, we labeled rifabutin orange in Table 3, indicating a potentially significant clinically interaction (34). This would provide insight into whether compatible regimens for cotreatment of HCV infection and drug-susceptible TB treatment are available. Additional DDIs with a potential for clinical significance include (i) the use of bedaquiline with DAAs that inhibit CYP enzymes and hence may increase bedaquiline concentrations and alter the concentrations of its metabolites and (ii) the use of isoniazid and/or clofazimine with DAAs that are in part eliminated by CYP3A enzymes given their weak inhibition of this enzyme system. However, given the high tolerability of oral DAAs and no clear correlation of toxicity with concentrations, it is unclear if combination treatment with either isoniazid or clofazimine would lead to higher rates of clinically significant adverse events (20). With regard to use of bedaquiline with DAAs, the potential clinical significance is predicated in large part on the concern for a narrow therapeutic window of bedaquiline, including an early study which showed a higher risk of death in patients receiving bedaquiline and known effects on QT prolongation (76). However, the relationship between bedaquiline concentrations and adverse events is unclear, and recent clinical data show bedaquiline is well tolerated (77). The lack of clinically significant DDIs between other anti-TB drugs used for MDR-TB and DAAs opens up the potential for cotreatment of



patients with MDR/extensively drug-resistant (XDR)-TB and HCV coinfection and highlights the need for pharmacokinetic studies evaluating DDIs. The recent WHO guidelines for MDR-TB list linezolid, fluoroquinolones, and bedaquiline as the key drugs (class A) for treatment, and given few predicted DDIs between DAAs and both linezolid and fluoroquinolones, DDI studies with DAA and bedaquiline should be an initial priority (16). If found that bedaquiline and some or all DAAs have limited or no clinically significant DDIs, this would pave the way for clinical cotreatment studies. In contrast to the Liverpool HEP online drug interaction tool, we listed the combination of delamanid (metabolites eliminated by CYP3A4) with any HCV drug which has inhibition of CYP enzymes as a potentially weak interaction unlikely to be clinically significant (Table 4).

Another important point to bring up with regard to potential DDIs between anti-TB and HCV infection treatment is the therapeutic window of oral DAAs. As detailed in the recent, comprehensive pharmacokinetics-pharmacodynamics (PK-PD) review of DAAs by Smolders et al. (20), the standard dose of most oral DAAs is at the higher end of doses tested given their high tolerability; however, with the exception of sofosbuvir (78), the antiviral effect was similar at lower doses. Thus, the clinical impact of anti-TB drugs which may decrease the concentrations of anti-HCV treatment such as the rifamycins is uncertain given DAA concentrations may still remain in an effective range.

In an effort to provide interim guidance to clinicians who decide to treat TB and HCV infection simultaneously, we provide the following recommendations. If treating a patient with drug-susceptible TB and HCV (genotypes 1 and 4 to 6) infection, we would favor the combination of isoniazid, rifabutin, pyrazinamide, and ethambutol with ledipasvir and sofosbuvir. The rationale being that there are fewer potential DDIs with ledipasvir and sofosbuvir than with other first-line HCV treatment regimens (Table 3). With regard to patients with drug-susceptible TB and infections with HCV genotypes 2 and 3, we would recommend the same anti-TB regimen as described above including rifabutin, given less potential for DDIs versus the other rifamycins and either glecaprevir-pibrentasvir or sofosbuvir-velpatasvir. Among patients with drug-resistant TB, we would recommend the newly recommended regimen (16) of three class A drugs (fluoroquinolone, bedaquiline, and linezolid) along with at least one class B drug (clofazimine or cycloserine) assuming susceptibility along with an HCV regimen of sofosbuvir and velpatasvir. This combination has the least theoretical DDIs (Table 4).

## LIVER DISEASE AND SUBSTANCE USE DISORDER

Although no TB drugs are recommended to be dose adjusted in patients with preexisting hepatic disease (Table 5), close and careful monitoring of liver function is needed, especially for unstable or advanced hepatic disease, as some studies have found such patients to be at higher risk of drug-related hepatotoxicity (79). Many TB drugs are metabolized in the liver; hence, the potential impact of liver disease severity on anti-TB drug pharmacokinetics should be addressed. While some data indicate that patients with chronic hepatitis, without cirrhosis, have comparable hepatic enzymatic activity to that of healthy individuals (80), there is also research finding moderate decreases in CYP enzyme activity in treatment-naïve HCV patients without significant liver disease (81). In patients with liver cirrhosis, significant impairment in hepatic clearance can occur, including decreases in the expression of some CYP enzymes (e.g., CYP3A4 and -1A2) and transporters (82). Wang et al. have shown that cirrhosis due to HCV infection has different effects on the protein expression of various transporters (83): increased protein expression in MATE-1 (multidrug and toxin extrusion protein 1) and decreased expression of OCT1 (organic cation transporter 1) and P-gp. This could have an impact on anti-TB and HCV drugs that are substrates to those transporters (Tables 1 and 2). In addition, if hypoalbuminemia related to liver disease is present, this could affect drugs metabolized by albumin such as delamanid, which is a highly protein-bound drug (84, 85); in fact, administering delamanid to patients with hypoalbuminemia has been associated with increased risk of QT prolongation, most likely due to high drug concentration. In cases where ascites is present, this would create an additional space for hydrophilic drugs to distribute, resulting in lower plasma drug

**TABLE 5** Characteristics of anti-tuberculosis drugs related to liver disease

Drug	Reference(s)	Protein binding (%)	Dosing adjustments for hepatic disease (14)	Associated with hepatotoxicity
<b>First-line drugs</b>				
Rifampin	79, 95–97	72–94	None; use with careful clinical and laboratory monitoring	Yes
Rifapentine	41, 79, 97, 98	97–99	None; use with careful clinical and laboratory monitoring	Yes
Isoniazid	79, 95, 97	0–34	None; use with caution in stable liver disease	Yes
Pyrazinamide	46, 79, 95, 97	0–7	None; use with careful clinical and laboratory monitoring.	Yes
			Contraindicated in severe hepatic impairment	
Ethambutol	79, 95	4–24	None	No
<b>Second-line drugs</b>				
Levofloxacin	99, 100	24–38	None	No <sup>a</sup>
Moxifloxacin	100–102	26–50	None; use with caution secondary to risk of QT prolongation	No <sup>a</sup>
Amikacin	103, 104	0–11	None	No
Kanamycin	103, 104	0	None	No
Capreomycin	104		None	No
Ethionamide	51, 101, 104	30	None; use with caution	Yes
Cycloserine	47, 104		None; use with caution in alcohol-related hepatitis	No
Linezolid	105, 106	31	None; not evaluated in severe hepatic impairment	No; a single case has been reported
Clofazimine	107		Not studied, use with caution, dose reduction may be warranted	No
Bedaquiline	108, 109	>99.9	None; use with caution in severe hepatic impairment (not studied)	Yes
Delamanid	60, 110	99.5	None; not recommended in moderate and severe liver impairments	No
<i>p</i> -Aminosalicylic acid	53, 104, 111	58–73	None; use with close laboratory and clinical monitoring	Yes

<sup>a</sup>Although levofloxacin and moxifloxacin were associated with increased risk of acute liver injury compared to clarithromycin (100), they are generally not considered hepatotoxic drugs.

concentrations (86). The effect of liver disease can also go beyond distribution and metabolism. For instance, drug absorption can be affected due to reduced first-pass metabolism, resulting in higher bioavailability in patients with cirrhosis, as has been observed with carvedilol and midazolam (86). Given all these variables, predicting the changes in drug kinetics in patients with liver disease is challenging and necessitates the need for specific studies in patients with TB and HCV infection to better understand the effect of the coinfection and cotreatment on drug concentrations, with the ultimate goal of optimizing therapy. It also is important to keep in mind the drug selection and dosing adjustment could be extended to drugs that are renally cleared, given that patients with HCV infection are at a higher risk of developing and accelerating the progression of kidney disease (87). The routes of elimination of each drug, including those with renal excretion, are summarized in Tables 1 and 2. Additionally, dosing recommendations for anti-TB drugs are included in recently released TB treatment guidelines (14). With regard to HCV treatment regimens, for patients with chronic kidney disease stages 1 to 3, there are no dosage adjustment recommendations for first-line regimens, whereas for chronic kidney disease stage 4 or 5, the AASLD/IDSA treatment guidelines recommend using either regular dose elbasvir-grazoprevir or glecaprevir-pibrentasvir, given sofosbuvir is renally excreted (17).

Metabolism of FDA-approved medication-assisted treatment (MAT) drugs used to treat substance use opiate disorder can be impacted by TB and HCV medications (88). Both methadone and buprenorphine are metabolized by CYP3A4 among other CYP enzymes. As a potent CYP3A4 inducer, rifampin has the potential to precipitate opiate withdrawal symptoms in patients receiving concomitant TB and opioid replacement therapy (89, 90). Rifabutin is also a CYP3A4 inducer but does not appear to cause a significant interaction with methadone, while studies with buprenorphine are lacking (91). Conversely, anti-TB and HCV drugs which have some inhibitory effect on CYP enzymes (Tables 1 and 2) may increase concentrations of methadone and buprenorphine, thus necessitating MAT dose adjustments. There are also few clinical data on DDIs between opiate replacement therapies and current HCV drugs. Simeprevir, paritaprevir, ombitasvir, ritonavir, and dasabuvir may have clinically significant drug interactions with buprenorphine and should be monitored closely (92). Given that the coepidemic of TB and HCV infection is linked in most cases by injection drug use, it will

be crucial to address substance use disorder and implement MAT and hence to understand the pharmacokinetic implications of combining TB, HCV, and medication-assisted treatments.

## CONCLUSION

Our review of available data on the potential DDIs between anti-TB and HCV treatment reveals a large gap in research addressing this topic. While knowledge of drug metabolism and existing data dim the prospects of combining any rifampin-based TB treatment with DAAs, our interpretation of drug metabolism highlights the potential compatibility of many anti-TB and HCV treatment regimens. Given the magnitude of the TB and HCV epidemics and their overlap, cotreatment could have a huge public health impact in many parts of the world, especially countries with well-developed national TB programs and limited existing capacity to scale up HCV treatment. Our review highlights possible anti-TB and HCV drug combinations to test in DDI studies and other pertinent PK issues to consider among patients with TB and HCV disease. Lastly, we hope this work helps to move the conversation and research agenda on addressing simultaneous TB and HCV infection forward.

## ACKNOWLEDGMENTS

The study was funded in part by the National Institutes of Health National Institute of Allergy and Infectious Diseases (K23AI103044, R21AI122001, R03AI139871 to R.R.K.) and the NIH Fogarty International Center (D43TW007124).

## REFERENCES

- WHO. 2018. Global tuberculosis report 2018. World Health Organization, Geneva, Switzerland.
- WHO. 2017. Global hepatitis report 2017. World Health Organization, Geneva, Switzerland.
- Altice FL, Azbel L, Stone J, Brooks-Pollock E, Smyrnov P, Dvoriak S, Taxman FS, El-Bassel N, Martin NK, Booth R, Stöver H, Dolan K, Vickerman P. 2016. The perfect storm: incarceration and the high-risk environment perpetuating transmission of HIV, hepatitis C virus, and tuberculosis in Eastern Europe and Central Asia. *Lancet* 388:1228–1248. [https://doi.org/10.1016/S0140-6736\(16\)30856-X](https://doi.org/10.1016/S0140-6736(16)30856-X).
- Lomtadze N, Kupreishvili L, Salakaia A, Vashakidze S, Sharvadze L, Kempker RR, Magee MJ, del Rio C, Blumberg HM. 2013. Hepatitis C virus co-infection increases the risk of anti-tuberculosis drug-induced hepatotoxicity among patients with pulmonary tuberculosis. *PLoS One* 8:e83892. <https://doi.org/10.1371/journal.pone.0083892>.
- Reis NR, Lopes CL, Teles SA, Matos MA, Carneiro MA, Marinho TA, Filho JA, Espirito-Santo MP, Lampe E, Martins RM. 2011. Hepatitis C virus infection in patients with tuberculosis in Central Brazil. *Int J Tuberc Lung Dis* 15:1397–1402. <https://doi.org/10.5588/ijtld.10.0636>.
- Chien JY, Huang RM, Wang JY, Ruan SY, Chien YJ, Yu CJ, Yang PC. 2010. Hepatitis C virus infection increases hepatitis risk during anti-tuberculosis treatment. *Int J Tuberc Lung Dis* 14:616–621.
- Kim WS, Lee SS, Lee CM, Kim HJ, Ha CY, Kim HJ, Kim TH, Jung WT, Lee OJ, Hong JW, You HS, Cho HC. 2016. Hepatitis C and not hepatitis B virus is a risk factor for anti-tuberculosis drug induced liver injury. *BMC Infect Dis* 16:50. <https://doi.org/10.1186/s12879-016-1344-2>.
- Nader LA, de Mattos AA, Picon PD, Bassanesi SL, De Mattos AZ, Pineiro Rodriguez M. 2010. Hepatotoxicity due to rifampicin, isoniazid and pyrazinamide in patients with tuberculosis: is anti-HCV a risk factor? *Ann Hepatol* 9:70–74. [https://doi.org/10.1016/S1665-2681\(19\)31682-5](https://doi.org/10.1016/S1665-2681(19)31682-5).
- Vermehren J, Park JS, Jacobson IM, Zeuzem S. 2018. Challenges and perspectives of direct antivirals for the treatment of hepatitis C virus infection. *J Hepatol* 69:1178–1187. <https://doi.org/10.1016/j.jhep.2018.07.002>.
- Golabi P, Sayiner M, Bush H, Gerber LH, Younossi ZM. 2017. Patient-reported outcomes and fatigue in patients with chronic hepatitis C infection. *Clin Liver Dis* 21:565–578. <https://doi.org/10.1016/j.cld.2017.03.011>.
- Suthar AB, Harries AD. 2015. A public health approach to hepatitis C control in low- and middle-income countries. *PLoS Med* 12:e1001795. <https://doi.org/10.1371/journal.pmed.1001795>.
- Tsai MC, Lin MC, Hung CH. 2009. Successful antiviral and antituberculosis treatment with pegylated interferon-alfa and ribavirin in a chronic hepatitis C patient with pulmonary tuberculosis. *J Formos Med Assoc* 108:746–750. [https://doi.org/10.1016/S0929-6646\(09\)60400-9](https://doi.org/10.1016/S0929-6646(09)60400-9).
- Mallikaarjun S, Wells C, Petersen C, Paccaly A, Shoaf SE, Patil S, Geiter L. 2016. Delamanid coadministered with antiretroviral drugs or antituberculosis drugs shows no clinically relevant drug-drug interactions in healthy subjects. *Antimicrob Agents Chemother* 60:5976–5985. <https://doi.org/10.1128/AAC.00509-16>.
- Nahid P, Dorman SE, Alipanah N, Barry PM, Brozek JL, Cattamanchi A, Chaisson LH, Chaisson RE, Daley CL, Grzemska M, Higashi JM, Ho CS, Hopewell PC, Keshavjee SA, Lienhardt C, Menzies R, Merrifield C, Narita M, O'Brien R, Peloquin CA, Raftery A, Saukkonen J, Schaaf HS, Sotgiu G, Starke JR, Migliori GB, Vernon A. 2016. Official American Thoracic Society/Centers for Disease Control and Prevention/Infectious Diseases Society of America Clinical Practice Guidelines: treatment of drug-susceptible tuberculosis. *Clin Infect Dis* 63:e147–e195. <https://doi.org/10.1093/cid/ciw376>.
- WHO. 2017. Guidelines for treatment of drug-susceptible tuberculosis and patient care, update 2017. World Health Organization, Geneva, Switzerland. License CC BY-NC-SA 3.0 IGO.
- WHO. 2018. Treatment guidelines for multidrug- and rifampin-resistant tuberculosis, 2018 update (pre-final text). World Health Organization, Geneva, Switzerland.
- AASLD-IDSA. 2019. Recommendations for testing, managing, and treating hepatitis C. <http://www.hcvguidelines.org>. Accessed 22 March 2019.
- Egelund EF, Alsultan A, Peloquin CA. 2015. Optimizing the clinical pharmacology of tuberculosis medications. *Clin Pharmacol Ther* 98:387–393. <https://doi.org/10.1002/cpt.180>.
- McConachie SM, Wilhelm SM, Kale-Pradhan PB. 2016. New direct-acting antivirals in hepatitis C therapy: a review of sofosbuvir, ledipasvir, daclatasvir, simeprevir, paritaprevir, ombitasvir and dasabuvir. *Expert Rev Clin Pharmacol* 9:287–302. <https://doi.org/10.1586/17512433.2016.1129272>.
- Smolders EJ, Jansen AME, Ter Horst PGJ, Rockstroh J, Back DJ, Burger DM. 2019. Viral hepatitis C therapy: pharmacokinetic and pharmacodynamic considerations: a 2019 update. *Clin Pharmacokinet* 58:1237. <https://doi.org/10.1007/s40262-019-00774-0>.
- Baciewicz AM, Chrisman CR, Finch CK, Self TH. 2013. Update on rifam-

- pin, rifabutin, and rifapentine drug interactions. *Curr Med Res Opin* 29:1–12. <https://doi.org/10.1185/03007995.2012.747952>.
22. Burman WJ, Gallicano K, Peloquin C. 2001. Comparative pharmacokinetics and pharmacodynamics of the rifamycin antibacterials. *Clin Pharmacokinet* 40:327–341. <https://doi.org/10.2165/00003088-200140050-00002>.
  23. Battaglia R, Pianezzola E, Salgarollo G, Zini G, Strolin Benedetti M. 1990. Absorption, disposition and preliminary metabolic pathway of 14C-rifabutin in animals and man. *J Antimicrob Chemother* 26:813–822. <https://doi.org/10.1093/jac/26.6.813>.
  24. Acocella G. 1978. Clinical pharmacokinetics of rifampicin. *Clin Pharmacokinet* 3:108–127. <https://doi.org/10.2165/00003088-197803020-00002>.
  25. Sanofi-Aventis. 2017. Priftrin (rifapentine), package insert. Sanofi-Aventis, Bridgewater, NJ.
  26. Acocella G, Nonis A, Perna G, Patane E, Gialdroni-Grassi G, Grassi C. 1988. Comparative bioavailability of isoniazid, rifampin, and pyrazinamide administered in free combination and in a fixed triple formulation designed for daily use in antituberculosis chemotherapy. II. Two-month, daily administration study. *Am Rev Respir Dis* 138:886–890. <https://doi.org/10.1164/ajrccm/138.4.886>.
  27. Alfarsi O, Alghamdi WA, Al-Shaer MH, Dooley KE, Peloquin CA. 2017. Rifampin vs. rifapentine: what is the preferred rifamycin for tuberculosis? *Expert Rev Clin Pharmacol* 10:1027–1036. <https://doi.org/10.1080/17512433.2017.1366311>.
  28. Song SH, Chang HE, Jun SH, Park KU, Lee JH, Lee EM, Song YH, Song J. 2013. Relationship between CES2 genetic variations and rifampicin metabolism. *J Antimicrob Chemother* 68:1281–1284. <https://doi.org/10.1093/jac/dkt036>.
  29. Burman WJ, Gallicano K, Peloquin C. 1999. Therapeutic implications of drug interactions in the treatment of human immunodeficiency virus-related tuberculosis. *Clin Infect Dis* 28:419–429. <https://doi.org/10.1086/515174>.
  30. Battaglia R, Salgarollo G, Zini G, Montesanti L, Strolin Benedetti M. 1991. Absorption, disposition, and urinary metabolism of 14C-rifabutin in rats. *Antimicrob Agents Chemother* 35:1391–1396. <https://doi.org/10.1128/aac.35.7.1391>.
  31. Skinner MH, Hsieh M, Torseth J, Pauloin D, Bhatia G, Harkonen S, Merigan TC, Blaschke TF. 1989. Pharmacokinetics of rifabutin. *Antimicrob Agents Chemother* 33:1237–1241. <https://doi.org/10.1128/aac.33.8.1237>.
  32. Nightingale SD, Cameron DW, Gordin FM, Sullam PM, Cohn DL, Chaisson RE, Eron LJ, Spati PD, Bihari B, Kaufman DL. 1993. Two controlled trials of rifabutin prophylaxis against *Mycobacterium avium* complex infection in AIDS. *N Engl J Med* 329:828–833. <https://doi.org/10.1056/NEJM199309163291202>.
  33. Zhaoxu L, Jingcheng T, Jinnan Z. 2008. Rifabutin autoinduction is caused by involvement of cytochrome P450 and cholinesterase. *Pharmazie* 63:156–159.
  34. Blaschke TF, Skinner MH. 1996. The clinical pharmacokinetics of rifabutin. *Clin Infect Dis* 22 Suppl 1:S15–S21. [https://doi.org/10.1093/clinids/22.Supplement\\_1.S15](https://doi.org/10.1093/clinids/22.Supplement_1.S15).
  35. Keung AC, Eller MG, Weir SJ. 1998. Single-dose pharmacokinetics of rifapentine in elderly men. *Pharm Res* 15:1286–1291. <https://doi.org/10.1023/a:1011960428896>.
  36. Dooley KE, Bliven-Sizemore EE, Weiner M, Lu Y, Nuermberger EL, Hubbard WC, Fuchs EJ, Melia MT, Burman WJ, Dorman SE. 2012. Safety and pharmacokinetics of escalating daily doses of the antituberculosis drug rifapentine in healthy volunteers. *Clin Pharmacol Ther* 91:881–888. <https://doi.org/10.1038/clpt.2011.323>.
  37. Savic RM, Lu Y, Bliven-Sizemore E, Weiner M, Nuermberger E, Burman W, Dorman SE, Dooley KE. 2014. Population pharmacokinetics of rifapentine and desacetyl rifapentine in healthy volunteers: nonlinearities in clearance and bioavailability. *Antimicrob Agents Chemother* 58:3035–3042. <https://doi.org/10.1128/AAC.01918-13>.
  38. Nakajima A, Fukami T, Kobayashi Y, Watanabe A, Nakajima M, Yokoi T. 2011. Human arylacetamide deacetylase is responsible for deacetylation of rifamycins: rifampicin, rifabutin, and rifapentine. *Biochem Pharmacol* 82:1747–1756. <https://doi.org/10.1016/j.bcp.2011.08.003>.
  39. Keung A, Reith K, Eller MG, McKenzie KA, Cheng L, Weir SJ. 1999. Enzyme induction observed in healthy volunteers after repeated administration of rifapentine and its lack of effect on steady-state rifapentine pharmacokinetics: part I. *Int J Tuberc Lung Dis* 3:426–436.
  40. Mitchison DA. 1992. The Garrod Lecture. Understanding the chemotherapy of tuberculosis—current problems. *J Antimicrob Chemother* 29:477–493. <https://doi.org/10.1093/jac/29.5.477>.
  41. Reith K, Keung A, Toren PC, Cheng L, Eller MG, Weir SJ. 1998. Disposition and metabolism of 14C-rifapentine in healthy volunteers. *Drug Metab Dispos* 26:732–738.
  42. Preziosi P. 2007. Isoniazid: metabolic aspects and toxicological correlates. *Curr Drug Metab* 8:839–851. <https://doi.org/10.2174/138920007782798216>.
  43. Ellard GA, Gammon PT. 1976. Pharmacokinetics of isoniazid metabolism in man. *J Pharmacokinet Biopharm* 4:83–113. <https://doi.org/10.1007/BF01086149>.
  44. Mitchell JR, Zimmerman HJ, Ishak KG, Thorgerirsson UP, Timbrell JA, Snodgrass WR, Nelson SD. 1976. Isoniazid liver injury: clinical spectrum, pathology, and probable pathogenesis. *Ann Intern Med* 84:181–192. <https://doi.org/10.7326/0003-4819-84-2-181>.
  45. Desta Z, Soukhova NV, Flockhart DA. 2001. Inhibition of cytochrome P450 (CYP450) isoforms by isoniazid: potent inhibition of CYP2C19 and CYP3A. *Antimicrob Agents Chemother* 45:382–392. <https://doi.org/10.1128/AAC.45.2.382-392.2001>.
  46. Stada Pharmaceuticals Inc. 2004. Pyrazinamide, package insert. Stada Pharmaceuticals Inc., Cranbury, NJ.
  47. Gumbo T. 2018. Chemotherapy of tuberculosis, *Mycobacterium avium* complex disease, and leprosy. In Brunton LL, Knollmann BC, Hilal-Dandan R (ed), Goodman & Gilman's the pharmacological basis of therapeutics. McGraw Hill Medical, New York, NY.
  48. Edwards CQ, Smith CR, Baughman KL, Rogers JF, Lietman PS. 1976. Concentrations of gentamicin and amikacin in human kidneys. *Antimicrob Agents Chemother* 9:925–927. <https://doi.org/10.1128/aac.9.6.925>.
  49. Janssen Pharmaceuticals Inc. 2016. Levaquin (levofloxacin), package insert. Janssen Pharmaceuticals Inc., Titusville, NJ.
  50. Fung HB, Kirschenbaum HL, Ojofeimi BO. 2001. Linezolid: an oxazolidinone antimicrobial agent. *Clin Ther* 23:356–391. [https://doi.org/10.1016/s0149-2918\(01\)80043-6](https://doi.org/10.1016/s0149-2918(01)80043-6).
  51. Wyeth Pharmaceuticals Inc. 2016. Trecator (ethionamide) (package insert). Wyeth Pharmaceuticals Inc., Philadelphia, PA.
  52. Nariman S. 1988. Adverse reactions to drugs used in the treatment of tuberculosis. *Adverse Drug React Acute Poisoning Rev* 7:207–227.
  53. Donald PR, Diacon AH. 2015. Para-aminosalicylic acid: the return of an old friend. *Lancet Infect Dis* 15:1091–1099. [https://doi.org/10.1016/S1473-3099\(15\)00263-7](https://doi.org/10.1016/S1473-3099(15)00263-7).
  54. Feng PC, Fenselau CC, Jacobson RR. 1981. Metabolism of clofazimine in leprosy patients. *Drug Metab Dispos* 9:521–524.
  55. Bayer Healthcare Pharmaceuticals. 2008. Avelox (moxifloxacin hydrochloride), package insert. Bayer Healthcare Pharmaceuticals, Wayne, PA.
  56. Brillault J, De Castro WV, Harnois T, Kitzis A, Olivier JC, Couet W. 2009. P-Glycoprotein-mediated transport of moxifloxacin in a Calu-3 lung epithelial cell model. *Antimicrob Agents Chemother* 53:1457–1462. <https://doi.org/10.1128/AAC.01253-08>.
  57. Te Brake LH, Russel FG, van den Heuvel JJ, de Knecht GJ, de Steenwinkel JE, Burger DM, Aarnoutse RE, Koenderink JB. 2016. Inhibitory potential of tuberculosis drugs on ATP-binding cassette drug transporters. *Tuberculosis (Edinb)* 96:150–157. <https://doi.org/10.1016/j.tube.2015.08.004>.
  58. Liu K, Li F, Lu J, Liu S, Dorko K, Xie W, Ma X. 2014. Bedaquiline metabolism: enzymes and novel metabolites. *Drug Metab Dispos* 42:863–866. <https://doi.org/10.1124/dmd.113.056119>.
  59. Shimokawa Y, Sasahara K, Koyama N, Kitano K, Shibata M, Yoda N, Umehara K. 2015. Metabolic mechanism of delamanid, a new antituberculosis drug, in human plasma. *Drug Metab Dispos* 43:1277–1283. <https://doi.org/10.1124/dmd.115.064550>.
  60. Sasahara K, Shimokawa Y, Hirao Y, Koyama N, Kitano K, Shibata M, Umehara K. 2015. Pharmacokinetics and metabolism of delamanid, a novel anti-tuberculosis drug, in animals and humans: importance of albumin metabolism *in vivo*. *Drug Metab Dispos* 43:1267–1276. <https://doi.org/10.1124/dmd.115.064527>.
  61. Steiner R, Raguž-Lučić N, Erceg D. 2017. Direct-acting antivirals (daas): drug-drug interactions (DDIs) in the treatment of hepatitis C virus (HCV). In Smolic M (ed), Update on hepatitis C. IntechOpen, London, UK. <https://doi.org/10.5772/intechopen.70788>.
  62. Denning J, Cornpropst M, Flach SD, Berrey MM, Symonds WT. 2013. Pharmacokinetics, safety, and tolerability of GS-9851, a nucleotide analog polymerase inhibitor for hepatitis C virus, following single ascending doses in healthy subjects. *Antimicrob Agents Chemother* 57:1201–1208. <https://doi.org/10.1128/AAC.01262-12>.
  63. Kirby BJ, Symonds WT, Kearney BP, Mathias AA. 2015. Pharmacokinetic, pharmacodynamic, and drug-interaction profile of the hepatitis C virus



- NS5B polymerase inhibitor sofosbuvir. *Clin Pharmacokinet* 54:677–690. <https://doi.org/10.1007/s40262-015-0261-7>.
64. Janssen Therapeutics. 2017. Olysio (simeprevir), package insert. Janssen Therapeutics, Titusville, NJ.
  65. MacBrayne CE, Kiser JJ. 2016. Pharmacologic considerations in the treatment of hepatitis C virus in persons with HIV. *Clin Infect Dis* 63 Suppl 1:S12–S23. <https://doi.org/10.1093/cid/ciw220>.
  66. Gilead Sciences Inc. 2014. Harvoni (ledipasvir and sofosbuvir), package insert. Gilead Sciences Inc., Foster City, CA.
  67. AbbVie. 2018. Viekira Pak (ombitasvir, paritaprevir, and ritonavir tablets; dasabuvir tablets), package insert. AbbVie. North Chicago, IL.
  68. Bristol-Myers Squibb. 2015. Daklinza (daclatasvir), package insert. Bristol-Myers Squibb, Princeton, NJ.
  69. Merck Sharp & Dohme Corp. 2016. Zepatier (elbasvir and grazoprevir), package insert. Merck Sharp & Dohme Corp., Whitehouse Station, NJ.
  70. AbbVie Inc. 2017. Mavyret (glecaprevir and pibrentasvir), package insert. AbbVie Inc., North Chicago, IL.
  71. Gilead Sciences Inc. 2017. Vosevi (sofosbuvir, velpatasvir, and voxilaprevir), package insert. Gilead Sciences Inc., Foster City, CA.
  72. Gilead Sciences Inc. 2017. Epclusa (sofosbuvir and velpatasvir), package insert. Gilead Sciences Inc., Foster City, CA.
  73. German P, Mathias A, Brainard D, Kearney BP. 2016. Clinical pharmacokinetics and pharmacodynamics of ledipasvir/sofosbuvir, a fixed-dose combination tablet for the treatment of hepatitis C. *Clin Pharmacokinet* 55:1337–1351. <https://doi.org/10.1007/s40262-016-0397-0>.
  74. Lawitz EJ, Gruener D, Hill JM, Marbury T, Moorehead L, Mathias A, Cheng G, Link JO, Wong KA, Mo H, McHutchison JG, Brainard DM. 2012. A phase 1, randomized, placebo-controlled, 3-day, dose-ranging study of GS-5885, an NS5A inhibitor, in patients with genotype 1 hepatitis C. *J Hepatol* 57:24–31. <https://doi.org/10.1016/j.jhep.2011.12.029>.
  75. Svensson EM, Dooley KE, Karlsson MO. 2014. Impact of lopinavir-ritonavir or nevirapine on bedaquiline exposures and potential implications for patients with tuberculosis-HIV coinfection. *Antimicrob Agents Chemother* 58:6406–6412. <https://doi.org/10.1128/AAC.03246-14>.
  76. Diacon AH, Pym A, Grobusch MP, de los Rios JM, Gotuzzo E, Vasilyeva I, Leimane V, Andries K, Bakare N, De Marez T, Haxaire-Theeuwes M, Louis N, Meyvisch P, De Paeppe E, van Heeswijk RP, Dannemann B, TMC207-C208 Study Group. 2014. Multidrug-resistant tuberculosis and culture conversion with bedaquiline. *N Engl J Med* 371:723–732. <https://doi.org/10.1056/NEJMoa1313865>.
  77. Nunn AJ, Phillips PPJ, Meredith SK, Chiang CY, Conradie F, Dalai D, van Deun A, Dat PT, Lan N, Master I, Mbrahtu T, Meressa D, Moodliar R, Ngubane N, Sanders K, Squire SB, Torrea G, Tsogt B, Rusen ID, Stream Study Collaborators. 2019. A trial of a shorter regimen for rifampin-resistant tuberculosis. *N Engl J Med* 380:1201. <https://doi.org/10.1056/NEJMoa1811867>.
  78. Lawitz E, Rodriguez-Torres M, Denning JM, Albanis E, Cornpropst M, Berrey MM, Symonds WT. 2013. Pharmacokinetics, pharmacodynamics, and tolerability of GS-9851, a nucleotide analog polymerase inhibitor, following multiple ascending doses in patients with chronic hepatitis C infection. *Antimicrob Agents Chemother* 57:1209–1217. <https://doi.org/10.1128/AAC.01263-12>.
  79. Saukkonen JJ, Cohn DL, Jasmer RM, Schenker S, Jereb JA, Nolan CM, Peloquin CA, Gordin FM, Nunes D, Strader DB, Bernardo J, Venkataraman R, Sterling TR, ATS (American Thoracic Society) Hepatotoxicity of Antituberculosis Therapy Subcommittee. 2006. An official ATS statement: hepatotoxicity of antituberculosis therapy. *Am J Respir Crit Care Med* 174:935–952. <https://doi.org/10.1164/rccm.200510-1666ST>.
  80. Palatini P, De Martin S. 2016. Pharmacokinetic drug interactions in liver disease: an update. *World J Gastroenterol* 22:1260–1278. <https://doi.org/10.3748/wjg.v22.i3.1260>.
  81. Morcos PN, Moreira SA, Brennan BJ, Blotner S, Shulman NS, Smith PF. 2013. Influence of chronic hepatitis C infection on cytochrome P450 3A4 activity using midazolam as an *in vivo* probe substrate. *Eur J Clin Pharmacol* 69:1777–1784. <https://doi.org/10.1007/s00228-013-1525-5>.
  82. Elbekai RH, Korashy HM, El-Kadi AO. 2004. The effect of liver cirrhosis on the regulation and expression of drug metabolizing enzymes. *Curr Drug Metab* 5:157–167. <https://doi.org/10.2174/1389200043489054>.
  83. Wang L, Collins C, Kelly EJ, Chu X, Ray AS, Salphati L, Xiao G, Lee C, Lai Y, Liao M, Mathias A, Evers R, Humphreys W, Hop CE, Kumer SC, Unadkat JD. 2016. Transporter expression in liver tissue from subjects with alcoholic or hepatitis C cirrhosis quantified by targeted quantitative proteomics. *Drug Metab Dispos* 44:1752–1758. <https://doi.org/10.1124/dmd.116.071050>.
  84. Morgan DJ, McLean AJ. 1995. Clinical pharmacokinetic and pharmacodynamic considerations in patients with liver disease. An update. *Clin Pharmacokinet* 29:370–391. <https://doi.org/10.2165/00003088-199529050-00005>.
  85. Uildemolins M, Roberts JA, Rello J, Paterson DL, Lipman J. 2011. The effects of hypoalbuminaemia on optimizing antibacterial dosing in critically ill patients. *Clin Pharmacokinet* 50:99–110. <https://doi.org/10.2165/11539220-000000000-00000>.
  86. Verbeeck RK. 2008. Pharmacokinetics and dosage adjustment in patients with hepatic dysfunction. *Eur J Clin Pharmacol* 64:1147–1161. <https://doi.org/10.1007/s00228-008-0553-z>.
  87. Park H, Adeyemi A, Henry L, Stepanova M, Younossi Z. 2015. A meta-analytic assessment of the risk of chronic kidney disease in patients with chronic hepatitis C virus infection. *J Viral Hepat* 22:897–905. <https://doi.org/10.1111/jvh.12413>.
  88. McCance-Katz EF, Sullivan LE, Nallani S. 2010. Drug interactions of clinical importance among the opioids, methadone and buprenorphine, and other frequently prescribed medications: a review. *Am J Addict* 19:4–16. <https://doi.org/10.1111/j.1521-0391.2009.00005.x>.
  89. Kreek MJ, Garfield JW, Gutjahr CL, Giusti LM. 1976. Rifampin-induced methadone withdrawal. *N Engl J Med* 294:1104–1106. <https://doi.org/10.1056/NEJM197605132942008>.
  90. McCance-Katz EF. 2011. Drug interactions associated with methadone, buprenorphine, cocaine, and HIV medications: implications for pregnant women. *Life Sci* 88:953–958. <https://doi.org/10.1016/j.lfs.2010.09.016>.
  91. Brown LS, Sawyer RC, Li R, Cobb MN, Colborn DC, Narang PK. 1996. Lack of a pharmacologic interaction between rifabutin and methadone in HIV-infected former injecting drug users. *Drug Alcohol Depend* 43:71–77. [https://doi.org/10.1016/s0376-8716\(97\)84352-9](https://doi.org/10.1016/s0376-8716(97)84352-9).
  92. Roncero C, Villegas JL, Martinez-Rebollar M, Buti M. 2018. The pharmacological interactions between direct-acting antivirals for the treatment of chronic hepatitis C and psychotropic drugs. *Expert Rev Clin Pharmacol* 11:999–1030. <https://doi.org/10.1080/17512433.2018.1519392>.
  93. O'Connor R, O'Sullivan JF, O'Kennedy R. 1995. The pharmacology, metabolism, and chemistry of clofazimine. *Drug Metab Rev* 27:591–614. <https://doi.org/10.3109/03602539508994208>.
  94. Dhiman RK, Saraswat VA, Rajekar H, Reddy C, Chawla YK. 2012. A guide to the management of tuberculosis in patients with chronic liver disease. *J Clin Exp Hepatol* 2:260–270. <https://doi.org/10.1016/j.jceh.2012.07.007>.
  95. Alghamdi WA, Al-Shaer MH, Peloquin CA. 2018. Protein binding of first-line antituberculosis drugs. *Antimicrob Agents Chemother* 62:e00641-18. <https://doi.org/10.1128/AAC.00641-18>.
  96. Te Brake LH, Ruslami R, Later-Nijland H, Mooren F, Teulen M, Apriani L, Koenderink JB, Russel FG, Burger DM, Alisjahbana B, Wieringa F, van Crevel R, Aarnoutse RE. 2015. Exposure to total and protein-unbound rifampin is not affected by malnutrition in Indonesian tuberculosis patients. *Antimicrob Agents Chemother* 59:3233–3239. <https://doi.org/10.1128/AAC.03485-14>.
  97. Sharma SK, Sharma A, Kadiravan T, Tharyan P. 2013. Rifamycins (rifampicin, rifabutin and rifapentine) compared to isoniazid for preventing tuberculosis in HIV-negative people at risk of active TB. *Cochrane Database Syst Rev* 2013:CD007545. <https://doi.org/10.1002/14651858.CD007545.pub2>.
  98. Egelund EF, Weiner M, Singh RP, Prihoda TJ, Gelfond JA, Derendorf H, MacKenzie WR, Peloquin CA. 2014. Protein binding of rifapentine and its 25-desacetyl metabolite in patients with pulmonary tuberculosis. *Antimicrob Agents Chemother* 58:4904–4910. <https://doi.org/10.1128/AAC.01730-13>.
  99. Fish DN, Chow AT. 1997. The clinical pharmacokinetics of levofloxacin. *Clin Pharmacokinet* 32:101–119. <https://doi.org/10.2165/00003088-199732020-00002>.
  100. Paterson JM, Mamdani MM, Manno M, Juurlink DN, Canadian Drug Safety and Effectiveness Research Network. 2012. Fluoroquinolone therapy and idiosyncratic acute liver injury: a population-based study. *CMAJ* 184:1565–1570. <https://doi.org/10.1503/cmaj.111823>.
  101. Budha NR, Lee RE, Meibohm B. 2008. Biopharmaceutics, pharmacokinetics and pharmacodynamics of antituberculosis drugs. *Curr Med Chem* 15:809–825. <https://doi.org/10.2174/092986708783955509>.
  102. Barth J, Jager D, Mundkowski R, Drewelow B, Welte T, Burkhardt O. 2008. Single- and multiple-dose pharmacokinetics of intravenous moxifloxacin in patients with severe hepatic impairment. *J Antimicrob Chemother* 62:575–578. <https://doi.org/10.1093/jac/dkn212>.

103. Kirby WM, Clarke JT, Libke RD, Regamey C. 1976. Clinical pharmacology of amikacin and kanamycin. *J Infect Dis* 134 Suppl:S312–S315. [https://doi.org/10.1093/infdis/135.supplement\\_2.s312](https://doi.org/10.1093/infdis/135.supplement_2.s312).
104. Blumberg HM, Burman WJ, Chaisson RE, Daley CL, Etkind SC, Friedman LN, Fujiwara P, Grzemska M, Hopewell PC, Iseman MD, Jasmer RM, Koppaka V, Menzies RI, O'Brien RJ, Reves RR, Reichman LB, Simone PM, Starke JR, Vernon AA, American Thoracic Society, Centers for Disease Control and Prevention and the Infectious Diseases Society. 2003. American Thoracic Society/Centers for Disease Control and Prevention/Infectious Diseases Society of America: treatment of tuberculosis. *Am J Respir Crit Care Med* 167:603–662. <https://doi.org/10.1164/rccm.167.4.603>.
105. De Bus L, Depuydt P, Libbrecht L, Vandekerckhove L, Nollet J, Benoit D, Vogelaers D, Van Vlierberghe H. 2010. Severe drug-induced liver injury associated with prolonged use of linezolid. *J Med Toxicol* 6:322–326. <https://doi.org/10.1007/s13181-010-0047-0>.
106. Anonymous. 2008. Handbook of anti-tuberculosis agents. Introduction. *Tuberculosis (Edinb)* 88:85–86. [https://doi.org/10.1016/S1472-9792\(08\)70002-7](https://doi.org/10.1016/S1472-9792(08)70002-7).
107. Cariello PF, Kwak EJ, Abdel-Massih RC, Silveira FP. 2015. Safety and tolerability of clofazimine as salvage therapy for atypical mycobacterial infection in solid organ transplant recipients. *Transpl Infect Dis* 17: 111–118. <https://doi.org/10.1111/tid.12340>.
108. Lakshmanan M, Xavier AS. 2013. Bedaquiline - the first ATP synthase inhibitor against multi drug resistant tuberculosis. *J Young Pharm* 5:112–115. <https://doi.org/10.1016/j.jyp.2013.12.002>.
109. Janssen Therapeutics. 2012. Sirturo (bedaquiline), package insert. Janssen Therapeutics, Titusville, NJ.
110. Gler MT, Skripconoka V, Sanchez-Garavito E, Xiao H, Cabrera-Rivero JL, Vargas-Vasquez DE, Gao M, Awad M, Park SK, Shim TS, Suh GY, Danilovits M, Ogata H, Kurve A, Chang J, Suzuki K, Tupasi T, Koh WJ, Seaworth B, Geiter LJ, Wells CD. 2012. Delamanid for multidrug-resistant pulmonary tuberculosis. *N Engl J Med* 366:2151–2160. <https://doi.org/10.1056/NEJMoa1112433>.
111. Iwainsky H. 1988. Mode of action, biotransformation and pharmacokinetics of antituberculosis drugs in animals and man. In Bartmann K (ed). *Antituberculosis drugs. Handbook of experimental pharmacology*, vol 84. Springer, Berlin, Germany. [https://doi.org/10.1007/978-3-642-72873-0\\_6](https://doi.org/10.1007/978-3-642-72873-0_6).